

Original Research Article

Nutritional evaluation of *Cucumis callosus* - an underutilized crop of semi arid region of Haryana

Bhawana* & Dr. Anita Kumari

Dept of Nutrition Biology, School of interdisciplinary and life sciences,
Central University of Haryana, Mehrangarh, Haryana

Abstract

Cucumis callosus, commonly known as wild melon or "kachri" in India, is a member of the **Cucurbitaceae** family. This study aims to evaluate the nutritional value of this underutilized crop. Kachri fruits vary in size, ranging from small to large, and are oval-shaped with a yellowish-green color. They start off sour and become sweet when ripe. These fruits are predominantly found in semiarid regions. The research findings indicate that the Mesocarp of *Cucumis callosus* contains a high protein content of 38.12% and an ash content of 9.02%. Recent studies have highlighted the health benefits of consuming wild edible plants, focusing on their antioxidant activity, which is essential for preventing and managing diseases. The antioxidant activity of *Cucumis callosus* ranges from $13.96 \pm 0.47\%$ to $31.68 \pm 2.66\%$. Additionally, the total phenol content (TPC) is measured at 931.5 μg GAE/g original weight and 683 μg GAE/g original weight.

Keywords: *Cucumis callosus*, kachri, mesocarp, DPPH, methanolic extract, physico-chemical, nutritive value

*Correspondence

Ms. Bhawana
bhawanayadav0709@gmail.com
Dept of Nutrition Biology, School of interdisciplinary and life sciences,
Central University of Haryana, Mehrangarh, Haryana

Received: 11 March 2024 - Accepted: 3 June 2024 - Published: 28 June 2024

Citation for the paper: Bhawana, Kumari A. Nutritional evaluation of *Cucumis callosus* - an underutilized crop of semi arid region of Haryana. International Journal of Nutrition & Lifestyle 2024; 4(2): 94 - 105. DOI: 10.69859/ijnl.2024.v4i2003



Copyright: © 2024 by the authors. Licensee: Research Thrive Publications. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Introduction

Cucumis callosus is a species of cucumber that belongs to the family cucurbitaceae. The plant produces edible fruits and is often used in traditional medicine for its various health benefits ^[1]. Although it is utilized in form of vegetables but ripe fruit is also consumed in India both at times of scarcity and on regular days. Vitamin C (19.99mg/100gm) is known to be present in *Cucumis* fruits ^[2].

Fruits and vegetables present a unique difficulty when drying because they are typically heat sensitive. Dehumidification must be done under carefully regulated settings. Due to the expensive cost and high level of skill required in the artificially drying system, sun drying is becoming more popular for preserving vegetables. However, in light of the present issues with micronutrient deficiencies, nutrient conservation is unquestionably essential ^[2].

Morphology

It is an annual plant that typically grows at a height of 40-80 cm creeping vine. It has small yellow flowers that are usually 2 cm in diameter. The fruit of it is edible and cheaper. Having leaves which are about 2-5 cm in different lobular structures ^[3]. The quality of *Cucumis callosus* depends on the effective rate of drying to minimize nutritional loss. The good quality dried *Cucumis callosus* is obtained to keep the slice thickness of 5 mm and it should be dried at 60°C temperature as shown in figure 1 ^[1].

Origin and Distribution: This is a native to south Asia, including Nepal, India, Bangladesh, and other parts of Asia, such as Thailand, China and Indonesia. While in India it is widely distributed across many states, including Rajasthan, Haryana, Uttar Pradesh, West Bengal, Assam and Karnataka ^[4].

According to reports, *Cucumis sativus* and *C. hystrix* are among the species that exhibit diversity ^[5]. A single variety, *Cucumis sativus* makes up the core gene pool of *Cucumis sativus*, while *Cucumis hystrix* makes up the secondary gene pool. Several researchers

have emphasized the need of collecting *Cucumis* species from India based on the genetic variability distribution and genetic erosion status [6] [7].

The genetic variation of the cucurbits in North Western India, particularly in the Aravali ranges, is rich, but it is under danger due to the spread of newly released types [8].

Significance of study

Cucumis callosus is a species which is wild and belongs to the Cucurbitaceae family. It was studied that vitamins and minerals are found in moderate amount in this fruit [9]. The secondary metabolites are present widely which are proven for therapeutic properties. Perishable crops above 20% promote food security all over world are dried to increase shelf life. It was studied that this fruit shows antioxidant properties and possess bioactive compounds [10].

In India the fruit and fruit powder is used in making curries and different items such as, pickles, dosa, chutney, vegetables, salad etc [11].

In this study the objectives were taken by considering the medicinal value of mesocarp of kachri fruit, the study entitled “**Nutrition evaluation of *Cucumis callosus* an underutilized crop of semiarid region of Haryana**”.

Objective

- To study the nutritional parameter of mesocarp of *Cucumis callosus*.
- To study the physiochemical, functional and antioxidant properties of mesocarp of *Cucumis callosus*

Research Methodology

Sample collection: Kachri fruit were taken from the local market of Mahendergarh, Haryana to conduct the study, which was used further to form the extract.

Extract Preparation

Processing: After collection of Kachri fruit, washing then cutting of the sample collected was done, to extract the mesocarp of the fruit. Mesocarp is then cut into 5 mm small slices. Further it was kept for 6 hours in hot air oven at 60°C temperature, in a petri plate for complete moisture loss. The dried mesocarp was grinded to form the powdered form and then stored in an air tight container in dark, to perform the experiments further as shown in figure 1.



Figure 1 Sample preparation from kachri fruit (*Cucumis callosus*)

Physical parameters of Kachri fruit

Length & Breadth: Length and breadth of the fruit was measured by vernier caliper scale.

Functional parameter of mesocarp of *Cucumis callosus*

Water absorption capacity: A clean empty falcon was taken. 1gm of sample (Ws) was mixed with 10ml of distilled water in that empty falcon was weighed down (W1). This was further mixed for 4-5 minutes on vortex. The sample was kept aside for 3-4 hours for diffusion. After that, for 25 minutes at the speed of 2900 revolution per minute on 25°C, the sample was centrifuged. Then the mesocarp sample was air dried inside hot air oven at 55°C. The supernatant was discarded and weight of the falcon is noted down (W2).

$$\text{WAC (\%)} = \frac{W2 - W1}{W_s} \times 100$$

Where, W1 = weight of the falcon + sample, W2 = weight of the falcon tube + sediment

Ws = weight of the sample

Oil absorption capacity: A clean empty falcon was taken. 1gm of sample (Ws) was mixed with 10ml of soyabean oil in that empty falcon was weighed down (W1). This was further mixed for 4-5 minutes on vortex. The sample was kept aside for 3-4 hours for diffusion. After that, for 25 minutes at the speed of 2900 revolution per minute on 25°C, the sample was centrifuged. Then the sample mesocarp was air dried inside hot air oven at 55°C. The supernatant was discarded and weight of the falcon is noted down (W2).

$$\text{OAC (\%)} = \frac{W2 - W1}{W_s} \times 100$$

Where, W1 = weight of the falcon tube + sample W2 = weight of the falcon tube + sediment,

Ws = weight of the sample

Emulsion activity: A clean empty falcon was taken. 1gm of sample was mixed with 10ml of distilled water and 10ml of soyabean oil in that empty falcon. After that, for 10 minutes

at the speed of 2000 revolution per minute on 25°C, the sample was centrifuged. Emulsion activity was further measured by the following formula.

$$\text{EA (\%)} = \frac{\text{Emulsified layer height}}{\text{Total content in falcon height}} \times 100$$

Emulsion stability: A clean empty falcon was taken. 1gm of mesocarp sample was diluted with 10ml of distilled water and 10ml of soyabean oil in that empty falcon. After that, for 10 minutes at the speed of 2000 revolution per minute on 25°C, the sample was centrifuged. Then for 30 minutes, the sample was kept in water bath at 80°C, cooled for 15 minutes under tap water. Centrifugation for 10 minutes at the speed of 2000 revolution per minute on 25°C was repeated again. Emulsion stability was further measured by the following formula.

$$\text{ES (\%)} = \frac{\text{Emulsified layer height}}{\text{Total content in falcon height}} \times 100$$

Proximate composition of mesocarp of *Cucumis callosus*: In this part of the study, moisture, ash, crude fibre, acid insoluble ash, protein and fat contents were estimated using standard methods of AOAC, 2000 ^[12]. Carbohydrate content was calculated by difference of other proximate components from 100.

Antioxidant activity (%Inhibition) of mesocarp of *Cucumis callosus*: It was estimated by DPPH method.

Bioactive compounds of mesocarp of *Cucumis callosus*: In this total phenolic compounds and total flavonoids were determined by Follin's Ceocalteu method.

Results and Discussion

The present study was directed to check the nutritional parameter and the physiochemical, functional and antioxidant properties of mesocarp of *Cucumis callosus*. Results have been divided under following subheadings -

1. Physical parameter of kachri fruit

Physical parameters Color, shape, length, breadth and weight of kachri fruit *Cucumis callosus* is presented in table 1.

Parameters	Mean value $\pm$ SD
Colour	Light green and yellowish with dark greenspots.
Shape	Round and oval
Length(cm)	1.5 ± 0.3
Breadth(cm)	4.5 ± 1.5
Weight(gm)	45 5

Table 1: Parameter of kachri fruit

2. **Yield percentage:** The yield percentage of Mesocarp powder is obtained from half kg of *Cucumis callosus* is 125.43 ± 5.6 gm.

3. Functional parameter of mesocarp of *Cucumis callosus*

From table 2, it is evident that the water absorption capacity i.e. weight of solvent (water) absorbed by mesocarp in saturated state over the dry mesocarp weight was $279.66 \pm 1.25\%$. Where the oil absorption capacity was $117 \pm 1.63\%$. Emulsion activity shown table 2 was $25.57 \pm 0.35\%$. Where emulsion stability was observed as $23.33 \pm 0.39\%$. The percentage decrease is may be due to hightemperature treatment of mesocarp sample.

Functional parameters	Mean value $\pm$ SD
Water absorption capacity (%)	279.66 $\pm$ 1.25
Oil absorption capacity (%)	117 $\pm$ 1.63
Emulsion activity (%)	25.57 $\pm$ 0.35
Emulsion stability (%)	23.33 0.39

Table 2: Functional parameter of mesocarp of *Cucumis callosus*

4. Proximate composition of mesocarp of *Cucumis callosus*

Proximate content of the mesocarp of *Cucumis callosus* was seen in table 3, in which it was observed that the **Moisture content** in the fresh mesocarp (*Cucumis callosus*) was $5.18 \pm 0.02\%$. Where the **Ash content** in the fresh mesocarp sample was noted as $9.02 \pm 0.1\%$. The crude fiber content was $8.01 \pm 0.1\%$, while the Acid insoluble ash was $0.9 \pm 0.1\%$. Protein content of the mesocarp sample was 38.12 ± 4.36 mg/1gm. The amount of **crude fat** present in the mesocarp (*Cucumis callosus*) sample was $24.32 \pm 0.01\%$ as shown in table 3 below -

Parameters	Mean value $\pm$ SD
Moisture (%)	5.18 ± 0.02
Ash (%)	9.02 ± 0.1
Crude Fiber (%)	8.01 ± 0.1
Acid insoluble ash (%)	0.9 ± 0.1
Protein (%)	38.12 ± 4.36
Fat (%)	24.32 ± 0.01
Carbohydrate (%)	32.68 ± 5.61

Table 4.3: Proximate composition of mesocarp of *Cucumis callosus*

5. Antioxidants as % inhibition of DPPH:

Antioxidants were noted as % inhibition of DPPH at different concentration rises from $13.96 \pm 0.47\%$ to $31.68 \pm 2.66\%$.





Figure 2: Scavenging activity (%inhibition of DPPH) of different concentration of methanolic extract of mesocarp (*Cucumis callosus*)

6. Bioactive compounds of mesocarp of *Cucumis callosus*

6.1 Total Phenolic content:

The extraction methods and solvents used dissolve the natural components present in the plants. These components can be categorized as either polar or non-polar. Phenolic compounds, due to their hydroxyl group, are more readily dissolved in polar organic solvents. Methanol was chosen as the extraction solvent as a result. Mesocarp (*Cucumis callosus*) has a total phenol contents (TPC) of 931.5 μg GAE/g original weight from figure 3.



6.2 Total flavonoid content:

Flavonoids are secondary metabolic products with antioxidant activity, and the amount and location of free OH groups determine how potent they are. Mesocarp (*Cucumis callosus*) has a total phenol contents (TPC) of 683 µg GAE/g original weight as shown in figure 4.




Figure 4: Standard value absorbance for TFC (µg/ml) in different concentration

Conclusion

The research reveals that the Mesocarp of *Cucumis callosus* contains a high protein content of 38.12% and an ash content of 9.02%. Recently, there has been increasing interest in the health benefits of consuming wild edible plants. Numerous studies have focused on the plant's antioxidant activity, which plays a crucial role in preventing and managing diseases. Antioxidants are compounds that can neutralize harmful molecules called reactive oxygen species in the body. These species are produced naturally during cellular metabolism but can increase due to factors like pollution, stress, and a poor diet. By consuming foods rich in antioxidants, one can counteract the negative effects of reactive oxygen species and lower the risk of diseases associated with oxidative stress. Evaluating the antioxidant activity of wild edible plants has shown that higher levels of phenolic and flavonoid compounds can serve as valuable sources of natural antioxidants.

These compounds possess potent antioxidant properties, making the consumption of wild plants a potential strategy for reducing oxidative stress and related diseases.

References

1. Gojiya DK* and Vyas DM, 2015. Studies on Effect of Slice Thickness and Temperature on Drying Kinetics of Kothimbda (*Cucumis Callosus*) and its Storage. *JFood Process Technol* 6(1). doi: 10.4172/2157-7110.1000406
2. Gadekar MS and Dodiya NS. Heterosis and combining ability analysis for yield and yield contributing traits in chickpea (*Cicer arietinum* L.). *Legume Research – An International Journal*. 2013; 36(5): 373 - 379
3. Nikita K. Gadekar*, Pragati S. Raut, Bhagyashree Mokale and Dr. Gajanan S. Sanap 2023. Review: on cucumis callocus. *World Journal of Pharmaceutical Research*. 12(5).doi: 10.20959/wjpr20235-27734
4. Kerje T and grum M. The origin of melon, *Cucumis melo*: A review of the literature. *Acta Horticulturae*. 2000; 510(510): 37 -44
5. Arora RK and Nayar ER. Wild relatives of crop plants in India. *National Bureau of Plant Genetic Resources*. 1984; vi, 90 p., [16]
6. Umesh Chandra RD and Kalloo G. Variability in pointed gourd (*Trichosanthes dioica* roxb.) germplasm collected from Eastern Uttar Pradesh. *Indian J plant Genetic Resources*. 1995; 8(1): 149 – 155
7. UmeshChandra and Koppar MN. Distribution, domestication and diversity of *Cucumis* species complex in Indian sub continent. *Indian J of Plant Genetic Resources*. 1992; 5(2): 37 – 48
8. Dwivedi AD, Gopal K. Biosynthesis of silver and gold nano particles using *chenopodium album* leaf extract. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*. 2010; 360 (1-3): 27 – 33

9. Lakshmi T, Devaraj E, Rajagopal V, Bhullar SK, Rajendran R. Acacia catechu ethanolic bark extract induces apoptosis in human oral squamous carcinoma cells. *Journal of Advanced Pharmaceutical Technology & Research*. 2017; 8(4): 143 – 149
10. Vishwkarma P, Singh P, Tripathi NN. Nutritional and antioxidant properties of wild edible macrofungi from North Eastern Uttar Pradesh, India. *NIScPR Online Periodicals Repository*. 2016; 15(1): 143 - 148.
11. Shivhare Y, Singh P, Gadekar R, Soni P. Botanicals as Antioxidants: A Renovate Review. *Research J of Pharmacognosy and Phytochemistry*. 2010; 2(4): 255 – 259
12. [AOAC] Assn. of Official Analytical Chemists. 2000. Coffee and tea. In: *Official methods of analysis*. 17th ed. Gaithersburg , Md. : AOAC